

Absolute Difference of Square Sum and Sum Mean Prime Labeling of Some Star Graphs

Sunoj B S¹, Mathew Varkey T K²

¹(Department of Mathematics, Government Polytechnic College, Attingal, Kerala, India
spalazhi@yahoo.com)

² (Department of Mathematics, TKM College of Engineering, Kollam, Kerala, India
mathewvarkeytk@gmail.com)

ABSTRACT

Absolute difference of square sum and sum mean prime labeling of a graph is the labeling of the vertices with $\{1, 2, \dots, p\}$ and the edges with absolute difference of the mean of the squares of the labels of the incident vertices and the sum of the labels of the incident vertices. The greatest common incidence number of a vertex (*gcin*) of degree greater than one is defined as the greatest common divisor of the labels of the incident edges. If the *gcin* of each vertex of degree greater than one is one, then the graph admits absolute difference of square sum and sum mean prime labeling. Here we investigate some star related graphs for absolute difference of square sum and sum mean prime labeling,

Keywords - Graph labeling, prime labeling, prime graphs, star, square sum.

I. INTRODUCTION

All graphs in this paper are finite and undirected. The symbol $V(G)$ and $E(G)$ denotes the vertex set and edge set of a graph G . The graph whose cardinality of the vertex set is called the order of G , denoted by p and the cardinality of the edge set is called the size of the graph G , denoted by q . A graph with p vertices and q edges is called a (p, q) - graph.

A graph labeling is an assignment of integers to the vertices or edges. Some basic notations and definitions are taken from [1],[2],[3] and [4]. Some basic concepts are taken from Frank Harary [1]. In this paper we investigated absolute difference of square sum and sum mean prime labeling of some star related graphs.

Definition: 1.1 Let G be a graph with p vertices and q edges. The greatest common incidence number (*gcin*) of a vertex of degree greater than or equal to 2, is the greatest common divisor (gcd) of the labels of the incident edges.

II. MAIN RESULTS

Definition 2.1 Let $G = (V(G), E(G))$ be a graph with p vertices and q edges. Define a bijection $f : V(G) \rightarrow \{1, 2, \dots, p\}$ by $f(v_i) = i$, for every i from 1 to p and define a 1-1 mapping $f_{adssmp}^* : E(G) \rightarrow$ set of natural numbers N by

$f_{adssmp}^*(uv) = \frac{1}{2} |f(u)^2 + f(v)^2 - \{f(u)+f(v)\}|$. The induced function f_{adssmp}^* is said to be an absolute difference of square sum and sum mean prime labeling, if for each vertex of degree at least 2, the *gcin* of the labels of the incident edges is 1.

Definition 2.2 A graph which admits absolute difference of square sum and sum mean prime labeling is called an absolute difference of square sum and sum mean prime graph.

Theorem 2.1 Star $K_{1,n}$ ($n > 2$) admits absolute difference of square sum and sum mean prime labeling.

Proof: Let $G = K_{1,n}$ and let v_1, v_2, \dots, v_{n+1} are the vertices of G .

Here $|V(G)| = n+1$ and $|E(G)| = n$.

Define a function $f : V \rightarrow \{1, 2, \dots, n+1\}$ by

$$f(v_i) = i, \quad i = 1, 2, \dots, n+1$$

Clearly f is a bijection.

For the vertex labeling f , the induced edge labeling

f_{adssmp}^* is defined as follows

$$f_{adssmp}^*(v_1 v_{i+1}) = \frac{i(i+1)}{2}, \quad i = 1, 2, \dots, n.$$

Clearly f_{adssmp}^* is an injection.

$$\begin{aligned} \text{gcin of } (v_1) &= \text{gcd of } \{f_{adssmp}^*(v_1 v_2), \\ &\quad f_{adssmp}^*(v_1 v_3)\} \\ &= \text{gcd of } \{1, 3\} \\ &= 1, \quad i = 1, 2, \dots, n-2. \end{aligned}$$

So, *gcin* of each vertex of degree greater than one is 1. Hence $K_{1,n}$, admits absolute difference of square sum and sum mean prime labeling.

Theorem 2.2 Bistar $B(m, n)$ ($m, n > 2$) admits absolute difference of square sum and sum mean prime labeling.

Proof: Let $G = B(m, n)$ and let $v_1, v_2, \dots, v_{m+n+2}$ are the vertices of G .

Here $|V(G)| = m+n+2$ and $|E(G)| = m+n+1$.

Define a function $f : V \rightarrow \{1,2,---,m+n+2\}$ by

$$f(v_i) = i, i = 1,2,---,m+n+2.$$

Clearly f is a bijection.

For the vertex labeling f , the induced edge labeling

f_{adssmp}^* is defined as follows

$$f_{adssmp}^*(v_{m+1} v_i) = \frac{m^2+m+i^2-i}{2} \quad i = 1,2,---,m.$$

$$f_{adssmp}^*(v_{m+1} v_{m+2}) = (m+1)^2.$$

$$f_{adssmp}^*(v_{m+2} v_{m+2+i}) = m^2+3m+mi + \frac{i^2+3i}{2}, \quad i = 1,2,---,n.$$

Clearly f_{adssmp}^* is an injection.

$$\begin{aligned} \text{gcin of } (v_{m+1}) &= \text{gcd of } \{f_{adssmp}^*(v_m v_{m+1}), \\ &\quad f_{adssmp}^*(v_{m+1} v_{m+2})\} \\ &= \text{gcd of } \{m^2, (m+1)^2\} = 1. \end{aligned}$$

$$\begin{aligned} \text{gcin of } (v_{m+2}) &= \text{gcd of} \\ &\{f_{adssmp}^*(v_{m+1} v_{m+2}), f_{adssmp}^*(v_{m+2} v_{m+3})\} \\ &= \text{gcd of } \{(m+1)^2, (m+2)^2\} = 1. \end{aligned}$$

So, **gcin** of each vertex of degree greater than one is 1.

Hence $B(m,n)$, admits absolute difference of square sum and sum mean prime labeling.

Theorem 2.3 Double graph of star $K_{1,n}$ ($n > 2$) admits absolute difference of square sum and sum mean prime labeling.

Proof: Let $G = D(K_{1,n})$ and let $v_1, v_2, ---, v_{2n+2}$ are the vertices of G .

Here $|V(G)| = 2n+2$ and $|E(G)| = 4n$.

Define a function $f : V \rightarrow \{1,2,---,2n+2\}$ by

$$f(v_i) = i, i = 1,2,---,2n+2.$$

Clearly f is a bijection.

For the vertex labeling f , the induced edge labeling

f_{adssmp}^* is defined as follows

$$f_{adssmp}^*(v_1 v_{i+2}) = \frac{i(i+3)}{2} + 1, \quad i = 1,2,---,n.$$

$$f_{adssmp}^*(v_2 v_{i+2}) = \frac{i(i+3)}{2} + 2, \quad i = 1,2,---,n.$$

$$f_{adssmp}^*(v_1 v_{n+i+2}) = \frac{(n+i+2)^2-n-i}{2} - 1, \quad i = 1,2,---,n.$$

$$f_{adssmp}^*(v_2 v_{n+i+2}) = \frac{(n+i+2)^2-n-i}{2}, \quad i = 1,2,---,n.$$

Clearly f_{adssmp}^* is an injection.

$$\begin{aligned} \text{gcin of } (v_2) &= \text{gcd of } \{f_{adssmp}^*(v_2 v_3), \\ &\quad f_{adssmp}^*(v_2 v_4)\} \\ &= \text{gcd of } \{4, 7\} = 1. \end{aligned}$$

$$\begin{aligned} \text{gcin of } (v_1) &= \text{gcd of } \{f_{adssmp}^*(v_1 v_5), \\ &\quad f_{adssmp}^*(v_1 v_3)\} \\ &= \text{gcd of } \{10, 3\} = 1. \end{aligned}$$

$$\begin{aligned} \text{gcin of } (v_{i+2}) &= \text{gcd of } \{f_{adssmp}^*(v_2 v_{i+2}), \\ &\quad f_{adssmp}^*(v_1 v_{i+2})\} \\ &= \text{gcd of } \left\{ \frac{i(i+3)}{2} + 2, \frac{i(i+3)}{2} + 1 \right\} \\ &= 1, \quad i = 1,2,---,n. \end{aligned}$$

$$\begin{aligned} \text{gcin of } (v_{n+i+2}) &= \text{gcd of } \{f_{adssmp}^*(v_2 v_{n+i+2}), \\ &\quad f_{adssmp}^*(v_1 v_{n+i+2})\} \\ &= \text{gcd of } \left\{ \frac{(n+i+2)^2-n-i}{2}, \right. \\ &\quad \left. \frac{(n+i+2)^2-n-i}{2} - 1 \right\} \\ &= 1, \quad i = 1,2,---,n. \end{aligned}$$

So, **gcin** of each vertex of degree greater than one is 1. Hence $D(K_{1,n})$, admits absolute difference of square sum and sum mean prime labeling.

Example 2.1 $G = D(K_{1,4})$




fig – 2.1

Theorem 2.4 Splitting graph of star $K_{1,n}$ ($n > 2$) admits absolute difference of square sum and sum mean prime labeling.

Proof: Let $G = S'(K_{1,n})$ and let $v_1, v_2, ---, v_{2n+2}$ are the vertices of G .

Here $|V(G)| = 2n+2$ and $|E(G)| = 3n$.

Define a function $f : V \rightarrow \{1,2,---,2n+2\}$ by

$$f(v_i) = i, i = 1,2,---,2n+2.$$

Clearly f is a bijection.

For the vertex labeling f , the induced edge labeling

f_{adssmp}^* is defined as follows

$$f_{adssmp}^*(v_1 v_{i+2}) = \frac{i(i+3)}{2} + 1, \quad i = 1,2,---,n.$$

$$f_{adssmp}^*(v_2 v_{i+2}) = \frac{i(i+3)}{2} + 2, \quad i = 1,2,---,n.$$

$$f_{adssmp}^*(v_1 v_{n+i+2}) = \frac{(n+i+2)^2-n-i}{2} - 1, \quad i = 1,2,---,n.$$

Clearly f_{adssmp}^* is an injection.

$$\begin{aligned} \text{gcin of } (v_{i+2}) &= \text{gcd of } \{f_{adssmp}^*(v_2 v_{i+2}), \\ &\quad f_{adssmp}^*(v_1 v_{i+2})\} \\ &= \text{gcd of } \left\{ \frac{i(i+3)}{2} + 2, \frac{i(i+3)}{2} + 1 \right\} \\ &= 1, \quad i = 1,2,---,n. \end{aligned}$$

So, *gcin* of each vertex of degree greater than one is 1. Hence $S'(K_{1,n})$, admits absolute difference of square sum and sum mean prime labeling.

Example 2.2 $G = S'(K_{1,4})$




fig – 2.2

Theorem 2.5 Let G be the tensor product of star $K_{1,n}$ and path P_2 ($n > 2$). G admits absolute difference of square sum and sum mean prime labeling.

Proof: Let $G = K_{1,n} \otimes P_2$ and let $v_1, v_2, \dots, v_{2n+2}$ are the vertices of G .

Here $|V(G)| = 2n+2$ and $|E(G)| = 2n$.

Define a function $f : V \rightarrow \{1, 2, \dots, 2n+2\}$ by $f(v_i) = i, i = 1, 2, \dots, 2n+2$.

Clearly f is a bijection.

For the vertex labeling f , the induced edge labeling f_{adssmp}^* is defined as follows

$$f_{adssmp}^*(v_1 v_{i+1}) = \frac{i(i+1)}{2}, \quad i = 1, 2, \dots, n.$$

$$f_{adssmp}^*(v_{n+2} v_{n+i+2}) = (n+2)(n+i+1) + \frac{i(i-1)}{2}, \quad i = 1, 2, \dots, n.$$

Clearly f_{adssmp}^* is an injection.

gcin of (v_1) = gcd of $\{f_{adssmp}^*(v_1 v_2), f_{adssmp}^*(v_1 v_3)\}$
 = gcd of $\{1, 3\} = 1$.

gcin of (v_2) = gcd of $\{f_{adssmp}^*(v_{n+2} v_{n+3}), f_{adssmp}^*(v_{n+2} v_{n+4})\}$
 = gcd of $\{(n+2)^2, n^2+5n+7\}$
 = gcd of $\{(n+2)^2, n+3\} = 1$.

So, *gcin* of each vertex of degree greater than one is 1. Hence $K_{1,n} \otimes P_2$, admits absolute difference of square sum and sum mean prime labeling.

Theorem 2.6 Duplicating the apex vertex of star $K_{1,n}$ ($n > 2$) admits absolute difference of square sum and sum mean prime labeling.

Proof: Let G be the graph and let v_1, v_2, \dots, v_{n+2} are the vertices of G .

Here $|V(G)| = n+2$ and $|E(G)| = 2n$.

Define a function $f : V \rightarrow \{1, 2, \dots, n+2\}$ by

$$f(v_i) = i, i = 1, 2, \dots, n+2.$$

Clearly f is a bijection.

For the vertex labeling f , the induced edge labeling f_{adssmp}^* is defined as follows

$$f_{adssmp}^*(v_1 v_{i+2}) = \frac{i(i+3)}{2} + 1, \quad i = 1, 2, \dots, n.$$

$$f_{adssmp}^*(v_2 v_{i+2}) = \frac{i(i+3)}{2} + 2, \quad i = 1, 2, \dots, n.$$

Clearly f_{adssmp}^* is an injection.

gcin of (v_2) = gcd of $\{f_{adssmp}^*(v_2 v_3), f_{adssmp}^*(v_2 v_4)\}$
 = gcd of $\{4, 7\} = 1$.

gcin of (v_1) = gcd of $\{f_{adssmp}^*(v_1 v_5), f_{adssmp}^*(v_1 v_3)\}$
 = gcd of $\{10, 3\} = 1$.

gcin of (v_{i+2}) = gcd of $\{f_{adssmp}^*(v_2 v_{i+2}), f_{adssmp}^*(v_1 v_{i+2})\}$
 = gcd of $\{\frac{i(i+3)}{2} + 2, \frac{i(i+3)}{2} + 1\}$
 = 1, $i = 1, 2, \dots, n$.

So, *gcin* of each vertex of degree greater than one is 1. Hence G , admits absolute difference of square sum and sum mean prime labeling.

Theorem 2.7 Duplicating the apex vertex of star $K_{1,n}$ by an edge ($n > 2$) admits absolute difference of square sum and sum mean prime labeling.

Proof: Let G be the graph and let v_1, v_2, \dots, v_{n+3} are the vertices of G .

Here $|V(G)| = n+3$ and $|E(G)| = n+3$.

Define a function $f : V \rightarrow \{1, 2, \dots, n+3\}$ by $f(v_i) = i, i = 1, 2, \dots, n+3$.

Clearly f is a bijection.

For the vertex labeling f , the induced edge labeling f_{adssmp}^* is defined as follows

$$f_{adssmp}^*(v_i v_{i+1}) = i^2, \quad i = 1, 2, 3.$$

$$f_{adssmp}^*(v_1 v_3) = 3, \quad i = 1, 2, \dots, n.$$

$$f_{adssmp}^*(v_3 v_{i+4}) = \frac{i(i+7)+18}{2}, \quad i = 1, 2, \dots, n-1.$$

Clearly f_{adssmp}^* is an injection.

gcin of (v_1) = gcd of $\{f_{adssmp}^*(v_1 v_2), f_{adssmp}^*(v_1 v_3)\}$
 = gcd of $\{1, 3\} = 1$.

gcin of (v_2) = gcd of $\{f_{adssmp}^*(v_1 v_2), f_{adssmp}^*(v_2 v_3)\}$
 = gcd of $\{1, 4\} = 1$.

gcin of (v_3) = gcd of $\{f_{adssmp}^*(v_2 v_3), f_{adssmp}^*(v_3 v_4)\}$
 = gcd of $\{4, 9\} = 1$.

So, *gcin* of each vertex of degree greater than one is 1.

Hence G , admits absolute difference of square sum and sum mean prime labeling.

Theorem 2.8 Let G be the graph obtained by joining each vertex of star $K_{1,n}$ to vertices of path P_2 by edges ($n > 2$) admits absolute difference of square sum and sum mean prime labeling.

Proof: Let G be the graph and let v_1, v_2, \dots, v_{n+2} are the vertices of G .

Here $|V(G)| = n+2$ and $|E(G)| = 2n+1$.

Define a function $f : V \rightarrow \{1, 2, \dots, n+2\}$ by

$$f(v_i) = i, \quad i = 1, 2, \dots, n+2.$$

Clearly f is a bijection.

For the vertex labeling f , the induced edge labeling

f_{adssmp}^* is defined as follows

$$f_{adssmp}^*(v_1 v_{i+2}) = \frac{i(i+3)}{2} + 1, \quad i = 1, 2, \dots, n.$$

$$f_{adssmp}^*(v_2 v_{i+2}) = \frac{i(i+3)}{2} + 2, \quad i = 1, 2, \dots, n.$$

$$f_{adssmp}^*(v_1 v_2) = 3.$$

Clearly f_{adssmp}^* is an injection.

$$\begin{aligned} \text{gcin of } (v_2) &= \text{gcd of } \{f_{adssmp}^*(v_1 v_2), \\ &\quad f_{adssmp}^*(v_2 v_3)\} \\ &= \text{gcd of } \{1, 4\} = 1. \end{aligned}$$

$$\begin{aligned} \text{gcin of } (v_1) &= \text{gcd of } \{f_{adssmp}^*(v_1 v_2), \\ &\quad f_{adssmp}^*(v_1 v_3)\} \\ &= \text{gcd of } \{1, 3\} = 1. \end{aligned}$$

$$\begin{aligned} \text{gcin of } (v_{i+2}) &= \text{gcd of } \{f_{adssmp}^*(v_2 v_{i+2}), \\ &\quad f_{adssmp}^*(v_1 v_{i+2})\} \\ &= \text{gcd of } \left\{ \frac{i(i+3)}{2} + 2, \frac{i(i+3)}{2} + 1 \right\} \\ &= 1, \quad i = 1, 2, \dots, n. \end{aligned}$$

So, **gcin** of each vertex of degree greater than one is 1.

Hence G , admits absolute difference of square sum and sum mean prime labeling.

REFERENCES

- [1] Apostol. Tom M, Introduction to Analytic Number Theory, Narosa, (1998).
- [2] F Harary, Graph Theory, Addison-Wesley, Reading, Mass, (1972)
- [3] Joseph A Gallian, A Dynamic Survey of Graph Labeling, The Electronic Journal of Combinatorics(2015), #DS6, Pages 1 – 389.
- [4] T K Mathew Varkey, Some Graph Theoretic Operations Associated with Graph Labelings, PhD Thesis, University of Kerala 2000.